

Problem Set 1

1. Regression with $MA(1)$ errors

Consider a regression model with $MA(1)$ disturbance term

$$\begin{aligned}y_t &= x_t' \beta + u_t \\ u_t &= \varepsilon_t + \theta \varepsilon_{t-1}, \quad t = 1, \dots, T\end{aligned}\tag{1.1}$$

where $\varepsilon_t \sim iid(0, \sigma^2)$ with $\varepsilon_0 = 0$ and x_t non-stochastic.

1. Derive an expression for the covariance matrix, $\sigma^2 \Omega$, of the vector of disturbances $u = (u_1, \dots, u_T)'$ in terms of θ .
2. By using $\varepsilon_t = u_t - \theta \varepsilon_{t-1}$ recursively, starting from $\varepsilon_0 = 0$, $\varepsilon_1 = u_1$, find the lower triangular matrix L such that $\varepsilon = Lu$ where $\varepsilon = (\varepsilon_1, \dots, \varepsilon_T)'$.
3. Assume θ is known. Noting that $Var(Lu) = \sigma^2 I$, determine a method for computing the best linear unbiased estimator of β which does not require the construction and inversion of Ω .

*© 2007 by Christian Julliard. This document may be reproduced for educational and research purposes, so long as the copies contain this notice and are retained for personal use or distributed free.

2. $F - GLS$ estimation of the regression model with $AR(1)$ errors

Consider the model

$$\begin{aligned}y_t &= x_t' \beta + u_t \\u_t &= \phi u_{t-1} + \varepsilon_t \quad t = 1, \dots, T\end{aligned}\tag{2.1}$$

where $|\phi| < 1$, $\varepsilon_t \sim iid(0, \sigma^2)$, x_t' and ε_t are process independent and

$$p \lim \frac{1}{T} \sum_{t=1}^T x_t x_t' = \Sigma_{xx}$$

and Σ_{xx} is non-singular.

1. You estimate (2.1) by OLS and obtain the residual \hat{u}_t . You estimate ϕ by

$$\hat{\phi} = \frac{\sum_{t=2}^T \hat{u}_t \hat{u}_{t-1}}{\sum_{t=2}^T \hat{u}_{t-1}^2}.$$

Assuming that $\hat{\beta}_{OLS}$ is consistent, show that $\hat{\phi}$ is a consistent estimator of ϕ .

3. Maximum Likelihood of the $ARMA(1,1)$

Consider the $ARMA(1,1)$ process

$$y_t = \phi y_{t-1} + \varepsilon_t + \theta \varepsilon_{t-1} \quad t = 1, \dots, T$$

where $|\phi| < 1$, $\varepsilon_t \sim iidN(0, \sigma^2)$

1. Assuming, $y_1 = \varepsilon_1 = 0$ write down the log likelihood.
2. Obtain the FOCs wrt ϕ and θ .
3. Obtain

$$\frac{1}{T} I(\psi)$$

where $\psi := [\phi, \theta, \sigma^2]'$.

4. In finite sample we can approximate the distribution of $(\hat{\psi} - \psi_0)$ as mean zero normal with variance depending on $I(\hat{\psi})$. How would you estimate the covariance matrix?