

# LTCC Course: Graph Theory 2019/20

## §4 Probabilistic Methods and Random Graphs

Peter Allen\*

3 February 2020

### References

Chapter 11 of Diestel and Chapter VII of Bollobás are good sources for most of what we will cover in this course. For those interested in exploring this topic further, there are some excellent textbooks devoted entirely to this subject. The classic texts on random graphs are:

- B. Bollobás, *Random Graphs*, Cambridge University Press, 2nd Edition (2001).
- S. Janson, T. Łuczak and A. Ruciński, *Random Graphs*, Wiley (2000).

The probabilistic method is treated in great detail (with many surprising applications) in:

- N. Alon and J. Spencer, *The Probabilistic Method*, Wiley, 3rd Edition (2008).

### Random Methods

Let's start with the following three innocent-looking questions. For each of them we assume that  $n$  is sufficiently large.

**Question 1.** *Is there an  $n$ -vertex graph with neither a clique nor an independent set of size  $5 \log n$ ?*

**Question 2.** *Is there an  $n$ -vertex graph without cycles of length less than 1000 but with chromatic number at least 10000?*

**Question 3.** *Is there an  $n$ -vertex graph with maximum degree at most  $n^{2/3}$ , but for every pair  $(U, V)$  of sets of vertices with  $|U|, |V| \geq n^{1/2}$  there is an edge from  $U$  to  $V$ ?*

The answer to each of these questions is **yes**, but how might one go about proving that? The natural first reaction on being confronted by questions like these is to try and *construct* graphs with the required combination of properties. In each case, one of the two required properties demands that the graph has rather few edges, and the other demands rather many. It's hard to strike a balance, and any attempt to base a construction around some nice structure seems doomed to failure<sup>1</sup>.

---

\*These notes are based on Graham Brightwell's notes for LTCC Graph Theory Courses in 2009-12. They were updated by Julia Böttcher in 2015.

<sup>1</sup>In fact there are clever constructions of 'pseudorandom' graphs which solve the last two questions, but these usually use deep theorems from algebraic geometry or number theory in their proofs — and they are still not good enough for the first question!

In fact, for each of the problems above, by far the best way to solve the problem is not to give an explicit construction at all, but instead to “construct” the graph “at random”, and show that, with positive probability, the random graph constructed has the required combination of properties. To illustrate this idea let us consider the first question.

*Affirmative answer to Question 1.* What we do is to build our random graph in the most naïve possible way. We take some (large) set of vertices, say  $V = [n] = \{1, \dots, n\}$ . Then, for each pair of vertices, independently, we toss a fair coin, and put an edge between the pair if we get a head.

Let’s calculate the probability that a particular set  $C$  of  $c$  vertices forms a clique. This means that each of the  $\binom{c}{2}$  coin-tosses corresponding to the pairs in the clique was a head, an event with probability  $2^{-\binom{c}{2}}$ .

Now, the *expected* number of cliques of size  $c$  is given by:

$$\binom{n}{c} 2^{-\binom{c}{2}} \leq \frac{n^c}{c!} 2^{-\binom{c}{2}} \leq \left( \frac{ne}{c 2^{(c-1)/2}} \right)^c.$$

If  $c \geq 2 \log_2 n$ , then  $n \leq 2^{c/2}$ , and so the expected number of  $c$ -cliques is at most  $(e\sqrt{2}/c)^c$ . For  $c \geq 5$ , this expectation is at most  $1/3$ . This implies that the probability that the random graph has a clique of size  $c$  is at most  $1/3$ . The same calculation shows that the probability that the graph has an independent set of size  $c$  is also at most  $1/3$ .

So, with probability at least  $1/3$ , the random graph on  $n \geq 5$  vertices has neither a clique nor an independent set of size as large as  $2 \log_2 n$ .  $\square$

*The idea: We can calculate expectation using ‘linearity of expectation’ and estimate probabilities from expectations.*

This delightfully simple argument was first given by Paul Erdős in 1947, and it is one of the first instances of the successful use of the *probabilistic method* in combinatorics. One can, and indeed Erdős did, recast this entire proof as a *counting argument*. Of the  $2^{\binom{n}{2}}$  graphs with vertex set  $[n]$ , the number of them in the class  $A_C$  with a clique or independent set on the set  $C$  of size  $c$  is  $\dots$ , and then effectively the same calculation shows that there are some graphs in none of the sets  $A_C$ .

The result we’ve proved is usually stated in terms of *Ramsey numbers*, a topic we’ll return to next week. It says that, for all  $c \in \mathbb{N}$ , if  $n < \frac{c}{e} 2^{(c-1)/2} \frac{1}{2^{1/c}}$  there is a graph on  $n$  vertices containing neither a clique nor an independent set of size  $c$ . This is exactly what it means to say that the *Ramsey number*  $R(c, c)$  is at least  $\frac{c}{e} 2^{(c-1)/2} (1 - o(1))$ . (Here the  $o(1)$  refers to a term that tends to zero as  $c \rightarrow \infty$ .)

## Sperner’s Theorem

This isn’t a result about graphs, but rather about partially ordered sets. However the people who prove such results tend also to be graph theorists.

Consider the Boolean cube  $Q^n$ . This is the collection of all  $2^n$  subsets of the set  $[n]$ , ordered by inclusion. An *antichain* or *Sperner family* in  $Q^n$  is a family of subsets of  $[n]$ , no one of which is contained in another. One way to construct a large antichain is to take the family of all subsets of  $[n]$  of size exactly  $\lfloor n/2 \rfloor$ . A famous theorem of Sperner from 1927 states that this is the largest antichain.

**Theorem 4** (Sperner’s theorem). *Antichains in  $Q^n$  have size at most  $\binom{n}{\lfloor n/2 \rfloor}$ .*

A stronger result is the so-called *LYM Inequality*.

**Theorem 5** (LYM inequality). *If  $\mathcal{A}$  is any antichain in  $Q^n$ , then*

$$\sum_{A \in \mathcal{A}} \binom{n}{|A|}^{-1} \leq 1.$$

This implies Sperner's Theorem, as  $\binom{n}{\lfloor n/2 \rfloor}$  is the largest binomial coefficient. Here's a snappy proof of the LYM Inequality.

*Proof.* Take any antichain  $\mathcal{A}$  in  $Q^n$ . Now take a random ordering  $(x_1, x_2, \dots, x_n)$  of  $[n]$ , and consider the *chain*  $\mathcal{C}$  of sets  $C_0, C_1, \dots, C_n$ , where  $C_j = \{x_1, \dots, x_j\}$ . Notice that, for each pair of elements of  $\mathcal{C}$ , one is contained in the other; thus  $\mathcal{C}$  intersects  $\mathcal{A}$  in at most one element. Hence it is certainly the case that  $\mathbb{E}|\mathcal{C} \cap \mathcal{A}| \leq 1$ .

Now, for each  $A \in \mathcal{A}$ , the probability that  $A$  is in the random chain  $\mathcal{C}$  is exactly  $1/\binom{n}{|A|}$ , since  $\mathcal{C}$  includes exactly one set with  $|A|$  elements, and it's equally likely to be any of them. So we have

$$1 \geq \mathbb{E}|\mathcal{A} \cap \mathcal{C}| = \sum_{A \in \mathcal{A}} \mathbb{P}(A \in \mathcal{C}) = \sum_{A \in \mathcal{A}} \frac{1}{\binom{n}{|A|}},$$

and we are done. □

*The idea: We find a clever random variable whose expectation is exactly the sum we want to bound, and whose expectation is 'obviously' at most one.*

Sperner's Theorem inspired a whole new line of research; a variety of generalisations are known by now. It also has interesting applications, among them the following result proved by Erdős in 1945 (this is also called the *Littlewood-Offord problem*; Littlewood and Offord came across this when studying the number of real roots of random polynomials).

**Theorem 6.** *Let  $x_1, \dots, x_n$  and  $y$  be real numbers with  $|x_i| \geq 1$  for each  $i$ . Then at most  $\binom{n}{\lfloor n/2 \rfloor}$  out of the  $2^n$  possible sums  $\sum_{i=1}^n \lambda_i x_i$  with  $\lambda_i \in \{-1, 1\}$  fall into the interval  $(y - 1, y + 1)$ .*

## Some probabilistic inequalities

So far we only used very simple probabilistic tools:

- the linearity of expectation,
- the fact that if the expectation of a (non-negative) random variable is smaller than one, then it takes the value zero with non-zero probability.

Before we can turn to more difficult results, we need to introduce some important probabilistic tools. Note that in these notes, and usually in probabilistic combinatorics in general, we will work with a finite probability space. This means that the 'usual' difficulties of measure theory disappear; we can assume all sets are measurable, we can replace integrals with finite sums and hence there is no question of convergence. If you aren't familiar with those terms: don't worry, the last two sentences just say you don't need to be.

**Lemma 7** (Markov's inequality). *Let  $X$  be any non-negative real-valued random variable. For each  $a > 0$ , we have  $\mathbb{P}(X \geq a) \leq \frac{\mathbb{E}X}{a}$ .*

*Proof.* By definition, we have

$$\mathbb{E}X = \sum_{i \in \mathbb{R}_0} i \mathbb{P}[X = i] \geq a \cdot \mathbb{P}[X \geq a]$$

which since  $a > 0$  completes the proof.  $\square$

Note that in the above proof, the ‘uncountably infinite’ sum is simply a technical convenience. Since  $X$  is a random variable on a finite probability space,  $X$  takes one of only finitely many values, and the sum is really a finite sum.

**Lemma 8** (Chebyshev’s inequality). *Let  $X$  be any real-valued random variable. For each  $k > 1$ , we have*

$$\mathbb{P}\left(|X - \mathbb{E}X|^2 \geq k^2(\mathbb{E}X^2 - (\mathbb{E}X)^2)\right) \leq \frac{1}{k^2}.$$

*Proof.* Observe that, letting  $Z = (X - \mathbb{E}X)^2$ , we have  $\mathbb{E}Z = \mathbb{E}X^2 - (\mathbb{E}X)^2$ , and  $Z$  is a non-negative random variable. Thus the desired statement is precisely Markov’s inequality applied to  $Z$ .  $\square$

For the third lemma we suppose that  $X_1, \dots, X_n$  are independent Bernoulli random variables, with  $\mathbb{P}(X_i = 1) = p_i$  (we do not need that the  $p_i$  are identical: but if they are we say  $X$  is a Binomial random variable). Let  $X = \sum_i X_i$ , so that by linearity of expectation we have  $\mathbb{E}X = \sum_i p_i$ .

**Lemma 9** (a Chernoff bound). *If  $X$  is the sum of independent Bernoulli random variables, then for each  $0 < \delta < 3/2$  we have*

$$\mathbb{P}(|X - \mathbb{E}X| \geq \delta \mathbb{E}X) < 2e^{-\frac{\delta^2 \mathbb{E}X}{3}}.$$

The proof of this is Exercise 1(a).

In all cases, the idea is supposed to be that a random variable is ‘usually’ ‘not too far’ from its mean. Markov’s inequality provides only weak bounds on what ‘usually’ and ‘not too far’ should be, but in return we have only to compute expectation. Chebyshev’s inequality often tells us much more, but we have to calculate not only the expectation of  $X$  but also of  $X^2$ , which can be tricky. Finally, Chernoff bounds (many different forms exist; the one quoted is often good) are very strong and easy to use, but only apply in the special case of a sum of independent Bernoulli random variables.

A very important fact, which we won’t use in these lectures but which is central to a lot of modern research, is that the last sentence is not really true. The Chernoff bound also applies to Bernoulli random variables which are *not* independent, but where the dependencies are ‘sequential’. For example, if we have  $\mathbb{P}(X_i = 1 | X_1, \dots, X_{i-1}) \leq p_i$  for each  $i$ , then letting  $S = \sum_i p_i$  we can conclude

$$\mathbb{P}(X \geq S + \delta S) < e^{-\frac{\delta^2 S}{3}}.$$

This can be proved using *coupling*. More excitingly, we can allow  $p_i$  to be not a constant but depend on the outcomes  $X_1, \dots, X_{i-1}$ . This means that  $S = \sum_i p_i$  is no longer a constant; it too is a random variable (the ‘observed expectation’). Suppose that we have  $S \leq s$  almost surely. Then we can still conclude

$$\mathbb{P}(X \geq (1 + \delta)s) < e^{-\frac{\delta^2 s}{3}}. \tag{1}$$

This is a *martingale concentration inequality* in disguise; proving it is Exercise 1(b). If you want to analyse a probabilistic process, you will likely need such things. A simple to state example is the

*triangle free process.* This starts with the graph  $G_0$  on  $n$  vertices with no edges. At each integer time  $t \geq 1$ , a pair  $(u, v)$  of vertices is chosen uniformly at random from the set of all pairs at distance three or more in  $G_{t-1}$  and the edge  $uv$  is added to form  $G_t$ . In other words, the constraint is that at each time we add an edge, we really add an edge ( $uv$  was not already present) and that edge does not create a triangle. The process stops when no such edges remain.

This process is easy to define, but hard to analyse (it was done in two independent papers, by Fiz Pontiveros, Griffiths and Morris, and by Bohman and Keevash, in 2013; both papers are quite long and technical). It turns out that the process is likely to run for about  $\frac{1}{2\sqrt{2}}n^{3/2}\sqrt{\log n}$  steps, and the final graph is likely to have very large independent sets (as triangle-free graphs on  $n$  vertices go); this gives the best lower bound we know on the Ramsey number  $R(3, k)$ —we’ll mention this again next week.

## Girth and Chromatic Number

Now let us return to Question 2, which we shall answer in a more general form in this section. The proof we shall give is yet another classic application of the probabilistic method, but one where we have to be more careful about our choice of the random object than before.

The *girth* of a graph  $G$  is the length of the shortest cycle in  $G$ . So a graph with high girth “locally” looks like a tree, hence has a rather simple structure. More precisely, suppose the girth is at least  $g$ , and we take  $2k < g$  and let  $N_k(v)$  be the number of vertices at distance at most  $k$  from vertex  $v$ . Then all the sets  $N_k(v)$  for  $v \in V(G)$  induce trees. In particular, all the induced subgraphs on the sets  $N_k(v)$  are 2-colourable. Is it nevertheless possible for the whole graph to have a large chromatic number? This is exactly what Question 2 asked for (for some specific parameters) and we will prove in this section that the answer is yes.

How can we prove that the chromatic number of a graph is large? One way is to show the graph has no large independent sets, as the set of vertices receiving any given colour is an independent set. Hence we will prove the following result.

**Theorem 10.** *For each  $g, k \in \mathbb{N}$  and all sufficiently large  $n$ , there exists a graph on  $n$  vertices with girth greater than  $g$  and no independent set of size at least  $\lceil n/k \rceil$ .*

The proof of this result (also due to Erdős, this time from 1959) is slightly more complicated than the one about Ramsey numbers. For a start, our random process involves the tossing of *biased* coins. Again we fix a vertex set  $[n]$ , but now we put an edge between each pair of vertices with probability  $p$ , all choices made independently. Here  $p = p(n)$  is a function of  $n$  that we can choose to suit our needs. What this defines is the standard (binomial) model  $\mathcal{G}(n, p)$  of *random graphs*. We use  $G(n, p)$  to denote a random graph chosen according to this method.

So our goal is to find some  $p$  such that (with positive probability, for sufficiently large  $n$ ) the random graph  $G(n, p)$  has neither an independent set of size at least  $n/k$ , nor a cycle of length at most  $g$ . Unfortunately there is no such  $p$ : if  $p = 100/n$ , it is very likely that  $G(n, p)$  contains triangles, and it is very likely that  $n/k$  vertices of  $G(n, p)$  are isolated (and in particular form an independent set) for some (large)  $k$ . So if  $p \leq 100/n$  we are likely to have too many isolated vertices, and if  $p > 100/n$  then we are likely to have triangles in  $G(n, p)$ . However, the idea then will be to choose the parameter  $p$  in such a way that  $G(n, p)$  ‘almost’ has the two properties we want so that a few vertex deletions will result in a graph with these properties.

*Proof.* Given  $k$  and  $g$  let  $\ell = \max\{k, g\}$  and take  $p(n) = n^{-1+\varepsilon}$ , where  $0 < \varepsilon < 1/\ell$ .

Let  $S \subset [n]$  be a fixed set of size  $\lceil n/\ell \rceil$ . How close is  $S$  to being independent in  $G(n, p)$ ? The number  $X$  of edges in  $S$  has expectation  $p \binom{|S|}{2}$ , and  $X$  is a random variable to which the Chernoff bound applies. In particular, we have

$$\mathbb{P}(|X| < \frac{pn^2}{8\ell^2}) < 2e^{-\frac{pn^2}{24\ell^2}} < e^{-n^{1+\varepsilon/2}} < \frac{1}{4}2^{-n}.$$

where the last two inequalities hold for all sufficiently large  $n$ . Intuitively, this says that it is unlikely that  $S$  is ‘near’ to being independent. There are at most  $2^n$  different choices for  $S \subset [n]$ . The probability that any one of them contains less than  $\frac{pn^2}{8\ell^2}$  edges in  $G(n, p)$  is therefore at most

$$2^n \cdot \frac{1}{4}2^{-n} = \frac{1}{4}.$$

So we conclude that for all sufficiently large  $n$ , with probability at least  $\frac{3}{4}$ , the random graph  $G(n, p)$  has the following property.

- (a) Every set  $S \subset [n]$  of size at least  $\lceil n/\ell \rceil$  contains at least  $\frac{pn^2}{8\ell^2} > n$  edges.

(Where the last inequality holds for sufficiently large  $n$ .)

Now we consider the expectation of  $N = N(G)$ , the number of cycles in  $G$  of length at most  $\ell$ .

$$\mathbb{E}N = \sum_{s=3}^{\ell} \frac{n(n-1) \cdots (n-s+1)}{2s} p^s \leq \frac{1}{2}(\ell-2)(np)^\ell.$$

Notice that  $(np)^\ell = n^{\varepsilon\ell}$ , so  $\mathbb{E}N = o(n)$ . Now,  $N$  is a non-negative random variable, and so by Markov’s inequality, with  $a = n/2$ , we find that  $\mathbb{P}(N \geq n/2) < 1/2$  for sufficiently large  $n$ . We conclude that with probability at least  $1/4$  the random graph  $G(n, p)$  has the following two properties.

- (a) Every set  $S \subset [n]$  of size at least  $\lceil n/\ell \rceil$  contains at least  $n$  edges,  
(b)  $N(G) < n/2$ .

This means that there is some graph  $H$  with both these properties. Fix such an  $H$ . So  $H$  has few cycles of length at most  $\ell$  and no set of size  $\lceil n/\ell \rceil$  is close to being independent.

In a last step we will now destroy these few cycles without deleting many edges in any of these sets. To this end, mark in each cycle of length at most  $\ell$  one edge (these edges need not be distinct) and delete the marked edges. Let the result be  $H'$ . Now we have deleted at most  $N(H) < n/2$  edges, so any set  $S \subset [n]$  of size at least  $\lceil n/\ell \rceil$  still contains at least  $n/2$  edges in  $H'$ . In other words,  $H'$  has no independent sets of size at least  $\lceil n/\ell \rceil$ , and it also has no cycles of length at most  $\ell$ .  $\square$

*The idea: For the  $p$  we chose,  $G(n, p)$  is ‘very far’ from having independent sets of size  $n/k$ , and we can use the Chernoff bound which gives us very strong probability bounds to check this. On the other hand, even though  $G(n, p)$  contains a few short cycles, it doesn’t have many. We can use Markov’s inequality to say that with a reasonable probability,  $G(n, p)$  has ‘not too many more than expectation’ short cycles, and we can delete them.*

With a similar strategy one can also given an affirmative answer to Question 3.

## Designs

Finally, we want to turn to a major classical problem in combinatorics, which was only solved very recently. Consider the following task: Given a complete graph on  $n$  vertices, partition the edges into as many edge-disjoint triangles as possible plus a left-over set. Is it possible to do this in such a way that the left-over set is empty?

A design theorist will recognise this as asking for a *Steiner triple system* on  $n$  elements. These are known to exist if and only if  $n$  is congruent to 1 or 3 modulo 6 (to see that this is necessary, observe that all vertices have to have even degree, so  $n$  must be odd, and the number of edges has to be divisible by 3, which rules out 5 modulo 6).

But what happens if we ask for edge-disjoint copies of  $K_k$  instead of  $K_3$  (and replaces “1 or 3 modulo 6” by appropriate other obvious divisibility conditions)? What happens if instead of partitioning the edges of a complete graph  $K_n$  we want to partition the edges of a complete  $t$ -uniform hypergraph  $K_n^{(t)}$  into copies of  $K_k^{(t)}$ ? These are all still good questions in design theory, but they are very hard. This problem dates back to Steiner in 1853, and it was a celebrated breakthrough when Keevash announced a solution at the beginning of 2014.

His solution, in fact, makes essential use of an *approximate* solution to the problem which was already given in 1985 by Rödl: He proved that the left-over set can be made an arbitrarily small fraction of the edge set (we do not need divisibility conditions in this case). To prove this approximate result Rödl used a (more advanced) probabilistic argument, which we shall discuss in the following.

**Theorem 11** (Rödl, 1985). *For any  $1 \leq t \leq k \in \mathbb{N}$  and  $\gamma > 0$ , if  $n$  is large enough, there is a partition of the edges of  $K_n^{(t)}$  into edge-disjoint copies of  $K_k^{(t)}$  and a leftover set of size at most  $\gamma n^t$ .*

To simplify the argument we will only consider the graph case for  $K_3$ -partitions here, that is, the case  $t = 2, k = 3, n > 3$ . Note that the  $t = 1$  case is trivial, as is the  $t = k$  case, as is the  $k = n$  case - so this is the smallest non-trivial case, but actually the general case is not much harder. So what we want to prove is that

- there is a set of  $(1 - \gamma)n^2/6$  edge-disjoint triangles in  $K_n$ .

Let us try a first approach at using the probabilistic method for this<sup>2</sup>: We have to somehow construct the partition ‘randomly’, and the obvious way is to select triangles from  $K_n$  ‘randomly’ in some way. So suppose we selected triangles from  $K_n$  independently at random with some probability  $p$ . Then the expected number of triangles we select is  $p\binom{n}{3}$ , so we should choose  $p$  to be at least  $(1 - \gamma)/n$  in order for the expected number of triangles to be at least  $(1 - \gamma)n^2/6$ . But it is easy to believe that these triangles will not be edge-disjoint: given any one triangle  $xyz$ , the expected number of (other) selected triangles containing  $xy$  is  $p(n - 3)$ , so the expected number of selected triangles which share an edge with  $xyz$  is  $3p(n - 3)$ , which is (at least) close to 3.

Maybe we can use a ‘trick’ like the one we used in proving Theorem 10, and delete some triangles in order to get a collection of edge-disjoint triangles? The only simple rule which will give us a collection of edge-disjoint triangles is: delete every triangle which shares an edge with another triangle. We can estimate how many selected triangles we have to delete by the number  $Y$  of pairs of selected triangles which share an edge. Hence  $\mathbb{E}Y = p^2\binom{n}{3} \cdot n \leq p^2n^4$ , and by Markov’s inequality we conclude that (with probability at least 9/10) we only have to delete  $10p^2n^4$  triangles. So we can find a collection of at least

$$p\binom{n}{3} - 10p^2n^4$$

---

<sup>2</sup>which will fail; then we will refine the method, which will also fail; and so on

edge-disjoint triangles. But unfortunately this is smaller than 0 if  $p \geq (1 - \gamma)/n$ . Even if we take  $p = \varepsilon/n$  for some  $\varepsilon > 0$ , this gives only approximately  $(\varepsilon - 60\varepsilon^2)\frac{n^2}{6}$  edge-disjoint triangles. This of course is (well) below  $n^2/6$ , so we cannot hope to even get close to the desired number of edge-disjoint triangles. But, if  $\varepsilon$  is very small then we will delete only a tiny proportion of the triangles we selected.

Rödl's clever insight now was the following. If we perform this select-and-delete method with  $p = \varepsilon/n$  for some very small  $\varepsilon > 0$ , then we will get a (small) collection of edge-disjoint triangles  $T_1$ , a (tiny) collection of 'waste' edges contained in the deleted triangles, and a (large) set of 'leftover' edges  $L_1$ . But this 'leftover' set will be very well-behaved - it will have the property that 'most' edges are contained in about the same number of triangles in  $L_1$ . We can repeat select-and-delete using only the edges of the leftover set to get another collection of edge-disjoint triangles  $T_2$  and another leftover set  $L_2$ . By construction  $T_1 \cup T_2$  is a collection of edge-disjoint triangles, and we can show that  $L_2$  is still well-behaved. We can keep taking these 'Rödl Nibbles' until we get down to a tiny leftover set and the desired collection of edge-disjoint triangles. That is the basic idea of the proof that follows.

Note that to justify that this procedure works, there are two things we need to show. First, we need to show that the leftover set is really always nice enough to run the 'nibble' procedure (and to analyse it!). Second, we need to show that the total number of 'waste' edges does not get large. If we show both of these, then when we stop we have by construction a collection of edge-disjoint triangles, and the edges not in that collection are either 'leftover' or 'waste'; both sets are small and therefore most edges are in the triangles, as we want. These two properties are established by the following lemma.

**Lemma 12.** *Given  $\gamma > 0$  and  $\varepsilon > 0$ , for all sufficiently small  $\delta' > 0$  there exists  $\delta > 0$  such that for all sufficiently large  $n$  the following holds. Suppose  $D \geq \gamma^2 n/32$ . Let  $G$  be a graph on  $n$  vertices in which all but at most  $\delta e(G)$  edges are in  $(1 \pm \delta)D$  triangles. Let  $S$  be a set of triangles selected independently with probability  $p = \varepsilon/D$  from the triangles of  $G$ . Let  $T$  be the set of triangles in  $S$  which share no edge with any other triangle of  $S$ . Let  $G'$  be obtained from  $G$  by deleting any edge contained in a triangle of  $S$ . Then with probability at least  $1/2$  we have*

$$(a) |S| = (1 \pm \delta')\varepsilon e(G)/3.$$

$$(b) |S \setminus T| < 60\varepsilon|S|.$$

$$(c) \text{ All but at most } \delta' e(G) \text{ edges of } G' \text{ lie in } (1 \pm \delta')(1 - \varepsilon)^2 D \text{ triangles of } G'.$$

We will only prove the following simplified version of this lemma. The difference is that the assumption on  $G$  that most of its edges are in about  $D$  triangles in Lemma 12 is replaced by the stronger assumption that all edges are in exactly  $D$  edges. (In 'real' we need the weaker assumption, because this is what (c) guarantees and we want to apply the lemma successively). Lemma 13 is not conceptually easier to prove than Lemma 12. But the stronger assumption makes the calculations substantially shorter.

**Lemma 13.** *Given  $\gamma > 0$  and  $\varepsilon > 0$ , for all sufficiently small  $\delta > 0$ , if  $n$  is large enough, the following holds. Suppose  $D \geq \gamma^2 n/32$ . Let  $G$  be a graph on  $n$  vertices in which every edge is in  $D$  triangles. Let  $S$  be a set of triangles selected independently with probability  $p = \varepsilon/D$  from the triangles of  $G$ . Let  $T$  be the set of triangles in  $S$  which share no edge with any other triangle of  $S$ . Let  $G'$  be obtained from  $G$  by deleting any edge contained in a triangle of  $S$ . Then with probability at least  $1/2$  we have*



(a)  $|S| = (1 \pm \delta)\varepsilon e(G)/3$ .

(b)  $|S \setminus T| < 30\varepsilon|S|$ .

(c) *All but at most  $\delta e(G)$  edges of  $G'$  lie in  $(1 \pm \delta)(1 - \varepsilon)^2 D$  triangles of  $G'$ .*

Before we prove Lemma 13 let us briefly (and sketchily) discuss how Lemma 12 implies Theorem 11. First, the bounds on  $|S|$  and  $|S \setminus T|$  yield bounds on  $e(G')$ : we find that  $e(G')$  is very close to  $(1 - \varepsilon)e(G)$  (the number of edges deleted is at least  $3|T|$  and at most  $3|S|$ ). We conclude that after  $t$  steps— $t$  nibbles—applying Lemma 12 starting from  $G_0 = K_n$  we get to a graph  $G_t$  in which the number of edges is close to  $(1 - \varepsilon)^t n^2/2$ , and most edges are in about  $(1 - \varepsilon)^{2t} n$  triangles. If we choose  $\tau$  such that  $\gamma/2 < (1 - \varepsilon)^\tau < \gamma$ , then we conclude that  $e(G_\tau) < \gamma n^2/2$  and that for each  $t \leq \tau$ , most edges of  $G_t$  are in at least  $(1 - \varepsilon)^{2t} n/2 > \gamma^2 n/8$  triangles. In particular this means the conditions of Lemma 12 that  $e(G)$  and  $D$  should not be too small are satisfied up to  $\tau$  steps, so we are allowed to make  $\tau$  nibbles.

Second, at each nibble step we partition the edges of  $G$  into three parts: the edges of  $T$  (which are edge-disjoint triangles), the edges in  $S \setminus T$  (which are ‘waste’), and the edges of  $G'$ . Because  $|S \setminus T|$  is much smaller than  $|T|$ , we conclude that the number of ‘waste’ edges in each nibble is a tiny fraction of the number of edges covered by  $T$ . So after  $\tau$  nibbles, we have partitioned  $E(K_n)$  into a collection of edge-disjoint triangles, a waste set which is tiny by comparison, and  $E(G_\tau)$  which we know has size at most  $\gamma n^2/2$ . Provided we chose  $\varepsilon$  sensibly, this means the waste set and  $E(G')$  together account for only at most  $\gamma n^2$  edges, and we conclude that the collection of edge-disjoint triangles we found covers all but  $\gamma n^2$  edges of  $E(K_n)$  as desired.

*Proof of Lemma 13.* Given  $\gamma > 0$  and  $\varepsilon > 0$ , we choose any sufficiently small  $\delta > 0$ .

Let  $G$  be an  $n$ -vertex graph in which every edge is in  $D$  triangles. Let  $S$  be obtained by choosing independently triangles from  $G$  with probability  $p = \varepsilon/D$ . Let  $T$  be the set of triangles in  $S$  which share no edge with any other triangle of  $S$ , and let  $G'$  be the graph obtained from  $G$  by removing all edges in triangles of  $S$ .

First we will estimate  $|S|$ . Observe that  $G$  has  $\frac{D}{3}e(G)$  triangles, and  $S$  is obtained by selecting independently from these triangles with probability  $p$ . So the expectation of  $|S|$  is  $\frac{D}{3}pe(G) = \varepsilon e(G)/3$ . By the Chernoff bound we conclude that

$$\Pr\left(|S| - \varepsilon e(G)/3 > \delta \varepsilon e(G)/3\right) < 2e^{-\delta^2 \varepsilon e(G)/3}$$

which (provided  $e(G)$  is large enough—which it is unless  $G$  is empty) is smaller than 0.1. We conclude that with probability at least 0.9 the set  $S$  has size

$$|S| = (1 \pm \delta)\varepsilon e(G)/3,$$

which proves that (a) holds with probability at least 0.9.

Now we can estimate  $|S \setminus T|$  by exactly the same method we tried to use above. We estimate how many pairs of triangles we find sharing an edge in  $S$ ; for each pair we delete two triangles. The number of pairs of triangles sharing an edge in  $G$  is just  $e(G)\binom{D}{2}$ , where we first choose the shared edge  $e$  then the pair of triangles. The expected number of these which appear in  $S$  is then  $p^2\binom{D}{2}e(G) < \varepsilon^2 e(G)/2$ . So the expected size of  $S \setminus T$  is at most  $\varepsilon^2 e(G)$  (since we delete both triangles from each pair). So by Markov’s inequality with probability at least 0.9 we have

$$|S \setminus T| \leq 10\varepsilon^2 e(G) < 60\varepsilon|S|,$$

which proves that (b) holds with probability at least 0.9.

Finally we have to show that most edges of  $G'$  are in the ‘right’ number of triangles of  $G'$ . This is the most difficult part of the proof. First, consider a fixed edge  $uv$  of  $G$ . We condition on the event that  $uv$  is also an edge of  $G'$ : that is, we assume no triangle containing  $uv$  is in  $S$ . We want to know  $T_{uv}$ , the number of triangles containing  $uv$  in  $G'$ . First we will try to find  $\mathbb{E}T_{uv}$ : for this we just need to find the probability that a given triangle  $uvw$  of  $G$  survives to  $G'$ . Now there are  $D - 1$  triangles of  $G$  (apart from  $uvw$ ) using the edge  $uw$ , and another (different!)  $D - 1$  triangles using the edge  $vw$ . The triangle  $uvw$  survives if and only if none of these triangles are in  $S$  ( $uvw$  cannot be in  $S$  since  $uv$  is an edge of  $G'$ ). The probability of that occurring is just  $(1 - p)^{2D-2}$ . So, since  $p = \frac{\varepsilon}{D}$ , we have

$$\mathbb{E}T_{uv} = D(1 - p)^{2D-2} \approx (1 - \varepsilon)^2 D.$$

Now we need to show that  $T_{uv}$  takes a value close to its expectation (at least most of the time). It would be very nice if we could say that  $T_{uv}$  was a sum of  $D$  independent Bernoulli random variables (one for each triangle of  $G$  containing  $uv$ ) and use the Chernoff bound, but this isn't true: these random variables are not independent, as it can happen that one triangle being selected affects two of the random variables. So we have to use Chebyshev's inequality, and this means we need to estimate  $\mathbb{E}T_{uv}^2$ .

**Claim 14.**

$$\mathbb{E}T_{uv}^2 \leq (\mathbb{E}T_{uv})^2 + \delta^5 D^2 / 2.$$

We will prove this later. Assuming the claim, we can complete the proof. By Chebyshev's inequality, we have

$$\mathbb{P}\left(|T_{uv} - \mathbb{E}T_{uv}|^2 \geq 2\delta^{-1}(\mathbb{E}T_{uv}^2 - (\mathbb{E}T_{uv})^2)\right) \leq \delta/16.$$

Substituting  $\mathbb{E}T_{uv}^2$ , we get that the probability of  $T_{uv} \neq \mathbb{E}T_{uv} \pm \delta^2 D$  is at most  $\delta/4$ . Since  $\delta < 1/4$ , and since  $D < 2\mathbb{E}T_{uv}$ , we conclude that the probability of  $T_{uv} \neq (1 \pm \delta/2)\mathbb{E}T_{uv}$  is at most  $\delta/4$ . Finally, since  $\mathbb{E}T_{uv}$  is approximately  $(1 - \varepsilon)^2 D$ , we conclude that the probability of  $T_{uv} \neq (1 \pm \delta)(1 - \varepsilon^2)D$  is at most  $\delta/4$ .

Now let  $Y$  be the random variable counting edges  $uv$  of  $G'$  for which  $T_{uv} \neq (1 \pm \delta)(1 - \varepsilon^2)D$ . The expectation of  $Y$  is at most  $\delta e(G)/4$ , so by Markov's inequality with probability at least  $3/4$  we have  $Y \leq \delta e(G)$ . In other words, with probability at least  $3/4$ , all but at most  $\delta e(G)$  edges of  $G'$  lie in  $(1 \pm \delta)(1 - \varepsilon)^2 D$  triangles of  $G'$ , which is (c).

Concluding, the probability that any one of (a), (b), (c) fails is at most  $0.1 + 0.1 + 0.25 < 0.5$ , so with probability at least  $0.5$  all three items hold as desired. It remains to check the Claim.

*Proof of Claim 14.* Given two events  $A$  and  $B$ , we define the *covariance* of  $A$  and  $B$  to be  $\text{Cov}(A, B) = \mathbb{E}AB - \mathbb{E}A\mathbb{E}B$ . Now if  $A$  and  $B$  are independent, their covariance is zero. This will help us evaluate  $\mathbb{E}T_{uv}^2$ .

For each  $w$  such that  $uvw$  is a triangle of  $G$ , let  $I_w$  be the random variable which takes value 1 if  $uvw$  is a triangle of  $G'$ , and 0 otherwise. We have the following equality.

$$\mathbb{E}T_{uv}^2 - (\mathbb{E}T_{uv})^2 = \mathbb{E}\left(\sum_w I_w\right)^2 - \left(\mathbb{E}\sum_w I_w\right)^2 = \mathbb{E}\sum_w I_w^2 + \mathbb{E}\sum_{w < w'} I_w I_{w'} - \sum_w (\mathbb{E}I_w)^2 - \sum_{w < w'} \mathbb{E}I_w \mathbb{E}I_{w'}.$$

and this is bounded above by

$$\mathbb{E}I_w^2 + \sum_{w < w'} \text{Cov}(I_w, I_{w'}) = \mathbb{E}T_{uv} + \sum_{w < w'} \text{Cov}(I_w, I_{w'}), \quad (2)$$

where we left out the third sum and rearranged. Our aim is to show that this last quantity is bounded above by  $\delta^5 D^2/2$ . Now consider some  $w$  and  $w'$  which make triangles of  $G$  with  $uv$ . The events  $I_w$  and  $I_{w'}$  are independent unless there is a triangle of  $G$  which uses both  $w, w'$  and one of  $u$  and  $v$ . In other words,  $\text{Cov}(I_w, I_{w'}) = 0$  unless either  $uww'$  or  $vww'$  (or both) are triangles of  $G$ . If one of the two (but not both) is a triangle of  $G$ , then

$$\text{Cov}(I_w, I_{w'}) = (1 - \varepsilon/D)^{4D-5} - (1 - \varepsilon/D)^{4D-4}$$

and if both are triangles, we get

$$\text{Cov}(I_w, I_{w'}) = (1 - \varepsilon/D)^{4D-6} - (1 - \varepsilon/D)^{4D-4}.$$

In either case, we get  $\text{Cov}(I_w, I_{w'}) \leq 2\varepsilon/D$ . Now for each given  $w$ , there are at most  $2D$  vertices  $w'$  such that one of  $uww'$  or  $vww'$  is a triangle of  $G$ . So (2) can be bounded above by

$$\mathbb{E}T_{uv} + \sum_w 2D(2\varepsilon/D) = \mathbb{E}T_{uv} + 4D < \delta^5 D^2/2,$$

where the final inequality is true since  $D > \gamma^2 n/16$ . □

That proves the Claim, and hence the proof of Lemma 13 is complete. □

## 1 Exercises

There are plenty more exercises in the textbooks!

**Exercise 1.** (a) Prove Lemma 9. Hint: first evaluate  $\mathbb{E}e^{tX}$  for  $t > 0$  a fixed constant, apply Markov's inequality, and optimise  $t$ . Start with the case of independent identical Bernoulli variables (i.e.  $p_1 = p_2 = \dots$ ). For the general case, you might want to look up 'Jensen's inequality'.

(b) Try to generalise this to prove the martingale concentration inequality (1). The difficult part is figuring out how to evaluate  $\mathbb{E}e^{tX}$ .

**Exercise 2.** For  $k \in \mathbb{N}$ , a graph  $G = (V, E)$  has Property  $S_k$  if, for every pair  $(A, B)$  of disjoint  $k$ -element subsets of  $V$ , there is a vertex  $x$  of the graph that is adjacent to every vertex of  $A$  and no vertex of  $B$ .

(a) Find a graph with property  $S_1$ .

(b) Show that, for each  $k \in \mathbb{N}$ , there is a graph with property  $S_k$ .

**Exercise 3.** A  $k$ -uniform hypergraph is a pair  $H = (V, E)$ , where  $V$  is a set of vertices, and  $E$  is a family of  $k$ -element subsets of  $V$ . (So a 2-uniform hypergraph is just a graph.) A hypergraph  $H = (V, E)$  has Property B if  $V$  can be partitioned into two subsets  $V_1$  and  $V_2$  in such a way that no edge is entirely contained within one of the two sets.

(a) Show that, if  $H = (V, E)$  is a  $k$ -uniform hypergraph with  $|E| < 2^{k-1}$ , then  $H$  has property B.

(b) Show that, if  $H = (V, E)$  is a  $k$ -uniform hypergraph such that each edge in  $E$  intersects at most  $d$  others, and  $e(d+1) \leq 2^{k-1}$ , then  $H$  has property B.

**Exercise 4.** (a) Let  $p = n^{-t}$ , for  $0 < t < 1$ , and let  $k$  be a fixed natural number. Write down an expression for the expected number of  $k$ -cliques in  $G(n, p)$ . Hence show that, if  $t > 2/(k-1)$ , the probability that  $G(n, p)$  contains a  $k$ -clique tends to zero as  $n \rightarrow \infty$ .

It is also true that, if  $t < 2/(k-1)$ , then the probability that  $G(n, p)$  contains a  $k$ -clique tends to one as  $n \rightarrow \infty$ : to prove this, one needs to work with the variance of the number of  $k$ -cliques.

(b) Let  $H$  denote the graph on five vertices  $a, b, c, d, e$  with seven edges:  $a, b, c, d$  form a clique, and  $de$  is also an edge. For  $p = n^{-7/10}$ , find the expected number of copies of  $H$  in  $G(n, p)$ . What is

$$\lim_{n \rightarrow \infty} \mathbb{P}(G(n, p) \text{ contains a copy of } H)?$$

(c) There is a parameter  $b(H)$  of graphs such that, if  $p = n^{-t}$  and  $t > b(H)$ , then the probability that  $G(n, p)$  contains a copy of  $H$  as a subgraph tends to zero, while if  $p = n^{-t}$  and  $t < b(H)$ , then this probability tends to 1. Based on the calculations in this question, what do you think this parameter  $b(H)$  might be?

**Exercise 5.** Set  $p = n^{-2/5}$ , and consider a random graph  $G = G(n, p)$ .

(a) Show that the degree of any fixed vertex  $v$  has a Binomial distribution, and find an upper bound on the probability that this degree is greater than or equal to  $n^{2/3}$ .

(b) Show that the probability that the maximum degree of  $G$  is at most  $n^{2/3}$  is at least  $2/3$ .

(c) Show that, with probability at least  $2/3$ , for every pair  $(U, V)$  of subsets of  $V(G)$ , with  $|U|, |V| \geq n^{1/2}$ , there is an edge from  $U$  to  $V$ .

(d) What can you deduce from (b) and (c)?

**Exercise 6.** (a) Try to prove Lemma 12. You should find that the only difficult part is to prove that most edges are in the ‘right’ number of triangles. You will not be able to prove that  $T_{uv}$  behaves nicely for every edge  $uv \in G$ : you will need to assume both that  $uv$  happened to lie in about the ‘right’ number of triangles in  $G$ , and that ‘most’ of those triangles share edges with about the ‘right’ number of triangles in  $G$ . If you make this assumption, you should be able to modify the argument given to show that  $T_{uv}$  is likely to be about the ‘right’ size. Then you will need to show that there cannot be too many edges of  $G$  which don’t satisfy the assumption.

(b) Try to prove the special case of Theorem 11 from Lemma 12. The difficulty here is to find out how to set constants in order to make the argument work.

(c) Try to prove Theorem 11—or try to understand the argument given in e.g. Alon and Spencer!