

LTCC Course: Graph Theory 2024/25

§5 Ramsey Theory and Regularity

Peter Allen & Leo Versteegen

17 March 2025

1 References

There is a wealth of material in Chapters 7 and 9 of Diestel, and a treasurehouse in Chapters IV and VII of Bollobás. Both books will also provide pointers to other sources.

2 Ramsey Theory

Ramsey theory is about results in the following style: no matter how “chaotic” the overall structure is, if we look at the right (usually small) piece of the structure, we will find a pattern. The most standard version of this type of result is the graph Ramsey theorem we saw in the previous lecture. In its most abstract form, Ramsey’s theorem says that, whatever finite number c of colours we are provided with, if n is large enough, then in every colouring of the edges of K_n with c colours, there is a monochromatic copy of K_k . We also quantify what size n is sufficient as follows.

Theorem 2.1. *If $n \geq 4^k$, then every 2-colouring of $E(K_n)$ contains a monochromatic copy of K_k .*

Proof. Let G be a two-coloured complete graph on $[n]$. We first construct a list of integers and colours a_1, \dots, a_{2k} and c_1, \dots, c_{2k} as follows. We let $a_1 = 1$ and c_1 be a majority colour on edges incident to a_1 (we break a tie arbitrarily). Now for each $2 \leq j \leq 2k$ sequentially, let S_j be the set of vertices joined to each a_i by colour c_i for $1 \leq i \leq j-1$. Let a_j be the smallest vertex of S_j , and c_j a majority colour on edges leaving a_j in S_j . It’s trivial to check $|S_j| \geq |S_{j-1}|/2$ for each j , so that we can construct the list. Now there is a majority colour among c_1, \dots, c_{2k} which corresponds to a monochromatic clique of size at least k among the a_1, \dots, a_{2k} . \square

The first result we saw last time shows that we cannot improve the condition $n \geq 4^k$ to (roughly) $n \geq \sqrt{2}^k$. But this is essentially all we know: we cannot prove either that 3.99^k is sufficient or that 1.42^k is not sufficient.

Another branch of Ramsey theory deals with colourings of the natural numbers and monochromatic sets which satisfy some ‘arithmetic’ condition. The original example is:

Theorem 2.2 (Schur, 1912). *For $c \in \mathbb{N}$, there exists $n = n(c) \in \mathbb{N}$ such that, for any c -colouring $f: [n] \rightarrow [c]$ of $[n]$, there are $x, y \in [n]$ such that x, y and $x + y$ all have the same colour.*

Proof. We choose n large enough so that any c -colouring of the edges of K_n contains a monochromatic triangle. Now, given $f: [n] \rightarrow [c]$, we construct a colouring g of the edges of K_n by the rule $g(ij) = f(|i - j|)$. By the choice of n there is a monochromatic triangle ijk with $i < j < k$. Now set $x = j - i$ and $y = k - j$. \square

This is the beginning of a rich theory. We say that the equation $z = x + y$ has a monochromatic solution in any c -colouring of $[n]$ for $[n]$ sufficiently large. Which equations, or systems of equations, have this property? This is the subject of Rado's partition calculus. (See Bollobás.)

A rather harder result is:

Theorem 2.3 (van der Waerden, 1927). *For each $c, r \in \mathbb{N}$ there exists $n = n(c, r)$ such that for any c -colouring of $[n]$ there exists a monochromatic arithmetic progression $a, a + d, \dots, a + (r - 1)d$ where $d \geq 1$.*

In yet another direction, there are geometric Ramsey statements.

Theorem 2.4 (Erdős and Szekeres, 1935). *For $k \in \mathbb{N}$ there exists $n = n(k) \in \mathbb{N}$ such that whenever X is a collection of n points in the plane in general position there is a set of k of the points that form the corners of a convex k -gon.*

Proof (sketch). Given a set X of points in the plane in general position we colour the 4-tuples of points "red" if the points form a convex 4-gon, and "blue" otherwise, i.e., if one of the points is inside the convex hull of the other three.

The hypergraph version of Ramsey's Theorem now says that we can either find k points such that all the 4-subsets are red – in which case we are done, as these points form a convex k -gon – or we can find k points such that all the 4-subsets are blue. But the latter is not possible, as there is no way to place even 5 points in the plane in general position without forming a convex 4-gon. \square

The theorems we saw so far all identify some "small part" of a structure which has a nice pattern. But there are also Ramsey results about larger substructures.

Theorem 2.5 (Gyárfás, 1983). *For any $n \in \mathbb{N}$ if the edges of K_n are coloured with red and blue, there exists a pair of vertex-disjoint paths which cover the vertices of K_n , one using only red edges and the other only blue edges.*

Observe that one of the paths in this result could be empty or only contain 1 vertex.

Proof. We construct the two paths P_1 and P_2 as follows. We start with both paths empty. Now for each $1 \leq j \leq n$ in succession we apply the following algorithm. If we can add j to the end of P_1 and maintain the property that P_1 uses only red edges, we do so. If not, but we can add j to P_2 and maintain the property that P_2 uses only blue edges, we do so. If we can do neither, then let u be an end of P_1 and v an end of P_2 . If uv is red, we remove v from P_2 and add v and j to P_1 , while if uv is blue we remove u from P_1 and add u and j to the end of P_2 . \square

3 Regularity

Szemerédi's regularity lemma has revolutionised graph theory. The purpose of this section is to give a short introduction to what the Lemma says, and how it is used. Let's start with a very loose and vague (and also false) statement of the regularity lemma. Basically, it says:

All graphs can be partitioned into a bounded number of vertex classes of the same size, so that the graph between any pair of classes resembles a random bipartite graph.

Let's try and make some sense of this. First of all, suppose B is a bipartite graph on the two equal vertex classes V_1 and V_2 (so every edge of B has one vertex in each class). What does it mean to say that “resembles a random bipartite graph”?

First of all, what is a random bipartite graph? We fix some $p \in [0, 1]$ and for each pair of vertices $u \in V_1, v \in V_2$ we put an edge between u and v with probability p , all choices made independently. Given $\varepsilon > 0$, using a Chernoff bound we can check that it is likely that (if $|V_1| = |V_2|$ is large enough) the following holds for every pair of sets $A \subset V_1$ and $B \subset V_2$ such that $|A| \geq \varepsilon|V_1|$ and $|B| \geq \varepsilon|V_2|$. The number of edges between A and B is $(p \pm \varepsilon)|A||B|$.

This motivates the following definition. We say a balanced bipartite graph B with partition classes V_1 and V_2 is ε -regular with density p if it “looks like” a random graph in this sense, that is, if for any pair of sets $A \subset V_1$ and $B \subset V_2$ of size $|A| \geq \varepsilon|V_1|$ and $|B| \geq \varepsilon|V_2|$, we have $(p \pm \varepsilon)|A||B|$ edges between A and B . It is convenient to reformulate this in terms of edge densities. Given a pair of disjoint vertex sets A and B the (edge) density of the pair (A, B) is

$$d(A, B) = \frac{e(A, B)}{|A||B|},$$

where $e(A, B)$ denotes the number of edges with one end in A and the other in B . If V_1, V_2 are disjoint vertex sets in a graph G and $d(V_1, V_2) = d$, then the pair (V_1, V_2) is ε -regular (in G) if the bipartite graph induced by V_1 and V_2 is ε -regular, which means that any pair of subsets $A \subset V_1$ and $B \subset V_2$ each covering at least an ε -fraction of V_1 and V_2 , respectively, has density $d(A, B) = d \pm \varepsilon$.

Now, let's return to our vague statement of the regularity lemma, and consider our demand that the partition of the vertex set into classes is such that “the graph between any pair of classes resembles a random bipartite graph”, or “every pair of classes is an ε -regular pair”. It turns out that this is too much to ask for (you'll see an example in the exercises). But we can replace “every” with “all but an ε -fraction of”.¹ What do we mean by saying that all the classes have the “same size”? Obviously we cannot mean this literally if, for example, the graph has a prime number of vertices. It turns out to be useful to allow ourselves to ignore a “small” set. This leads to the following definition.

Definition 3.1. For a graph G an ε -regular partition of G is a partition of $V(G)$ into classes V_0, V_1, \dots, V_k with $|V_0| \leq \varepsilon|V(G)|$ and $|V_1| = |V_2| = \dots = |V_k|$, such that all but at most $\varepsilon \binom{k}{2}$ pairs (V_i, V_j) are ε -regular. The set V_0 is also called the exceptional set.

Finally, we wanted a “bounded number” of vertex classes in our partition. Let us explain why. It is easy to check that for any graph G , the partition of $V(G)$ into singletons is a partition in which every pair of parts is ε -regular (for any ε); but this partition clearly is not useful. To exclude this (at least for large graphs), we impose an upper bound on the number of classes. This upper bound *can*, and certainly does, depend on ε , but it does not depend on the number of vertices of G . Another “useless partition” is the trivial partition which puts all vertices of G into the same class. To exclude this and more generally allow for a reasonably large number of classes we also want to impose a lower bound on their number.

Now we have made all parts of our claim above precise, which brings us to the following version of the regularity lemma.

Lemma 3.2 (Szemerédi's regularity lemma). For every $\varepsilon > 0$ and $k_0 \in \mathbb{N}$ there exists $K \in \mathbb{N}$ such that every graph G has an ε -regular partition into k classes with $k_0 \leq k \leq K$.

¹This ε is the same ε as in ε -regular. This is not necessary: we could have one error parameter for ε -regular and a different error parameter for the fraction of non-regular pairs. But we would not gain anything by doing this, so it simplifies notation to have only one error parameter.

Note that this is trivial for graphs with less than K vertices, because we will get the partition into singletons. But for graphs on much more than K vertices, this is a very strong statement. When we use the regularity lemma, we are usually trying to prove a result for graphs on n vertices, for all sufficiently large n , in which case we can think of each partition class as being a big set of vertices.

Also, the regularity lemma says nothing about what happens inside any one of the classes. This is one of the reasons why it is useful to be able to set a lower bound on the number of classes, so that the total number of edges inside the classes is guaranteed to be relatively small (at most $\varepsilon \binom{n}{2}$). We will only sketch the proof of this lemma here. For details (of this approach), see Diestel or Bollobás.

Proof of Lemma 3.2 (sketch). The basic idea of the proof is that we will describe an algorithm which starts with an initial partition into k_0 classes (we only need to ask that the classes are all close to the same size) and then successively refines the partition, at each step partitioning each part into a bounded number of new parts. Then we need to show that the algorithm terminates after a bounded number of steps. To do this we shall use a parameter called *mean square density* or *energy*, which we will now define. Given a partition \mathcal{P} of the vertices of the n -vertex graph G , we define

$$\mathcal{E}(\mathcal{P}) = \sum_{X \neq Y \in \mathcal{P}} d(X, Y)^2 |X||Y|/n^2.$$

This parameter behaves a little like the second moment $\mathbb{E}A^2$ of a random variable A : it measures the extent to which the densities between pairs of parts in \mathcal{P} fluctuate. (Exercise: in fact it is exactly the second moment of some random variable—which one?) It has two properties we need. Firstly,

$$\text{if } \mathcal{P}' \text{ refines } \mathcal{P}, \text{ then } \mathcal{E}(\mathcal{P}') \geq \mathcal{E}(\mathcal{P}).$$

This follows from a fairly easy application of the Cauchy-Schwarz inequality. Secondly, suppose (V_1, V_2) is not ε -regular. By definition this means that there are sets W_1 and W_2 of sizes at least $\varepsilon|V_1|$ and $\varepsilon|V_2|$ such that $d(W_1, W_2) \neq d(V_1, V_2) \pm \varepsilon$. Now the contribution of (V_1, V_2) to $\mathcal{E}(\mathcal{P})$ is $d(V_1, V_2)^2 |V_1||V_2|/n^2$. Hence, if we refine \mathcal{P} by splitting V_1 into W_1 and $V_1 \setminus W_1$, and V_2 similarly, we can easily check that

$$\text{the energy increases by at least } \varepsilon^4 |V_1||V_2|/n^2.$$

This is an easy calculation.

So suppose \mathcal{P} is a partition in which all parts are (approximately) the same size, but it is not ε -regular. Then we can imagine performing one refinement as above for each pair which is not ε -regular to get \mathcal{P}' . In this way² we split each class of \mathcal{P} into at most $2^{|\mathcal{P}|-1}$ classes. We get an increase in energy of about $\varepsilon^4/(|\mathcal{P}|^2)$ for each pair we refined, of which there are at least $\varepsilon \binom{|\mathcal{P}|}{2}$. So we will get a total increase in energy of at least $\varepsilon^5/4$. Of course \mathcal{P}' may well have parts of very different sizes. But using the first property we can refine \mathcal{P} further to a partition \mathcal{P}'' in which all classes are about the same size. We have

$$\mathcal{E}(\mathcal{P}) + \varepsilon^5/4 \leq \mathcal{E}(\mathcal{P}') \leq \mathcal{E}(\mathcal{P}'')$$

where the second inequality follows from the first property above. Now either \mathcal{P}'' is an ε -regular partition and we are done, or we can repeat the refinement.

However, we cannot repeat the refinement more than $4\varepsilon^{-5}$ times, otherwise we would reach a partition whose energy is larger than one, which is impossible as we always have $d(V_i, V_j) \leq 1$. \square

²In fact we need to be a little careful in this process so that we do not create very small clusters; we can deal with this by ‘merging’ any such very small clusters.

We will see how and why this is a useful result in the next section. But here is a note of caution. We would like to apply the result with a reasonably small value of ε , but then how large does $K = K(\varepsilon)$ have to be?³ This proof of the Regularity Lemma shows that we can take K to be a tower of twos of *height* at most $4\varepsilon^{-5}$. Moreover, Gowers showed that this is really something like the truth: a tower of twos of height at least $\varepsilon^{-1/16}$ is needed! So as soon as we prove something using the Regularity Lemma, we introduce unpleasantly large constants.

There are a lot of reasonable questions one can ask about the Regularity Lemma and its proof. Why did we take mean square density? The answer is that we could actually work with any strictly convex function (replacing the Cauchy-Schwarz inequality with Jensen’s inequality), but the calculations are easiest with this function. Is it important that we actually never referred to single edges but only edge densities? The answer to this is that it is definitely important: the whole proof goes through taking (measurable) partitions of $[0, 1]$ to deduce “structure” for any (measurable) $f: [0, 1]^2 \rightarrow I$, where I can be any bounded interval. This leads to a recent topic in combinatorics called “Graph Limits”. Is there any “other” proof of the Regularity Lemma? There is: one can show that, given $\varepsilon > 0$, if (a very large but bounded number of) vertices v_1, \dots, v_ℓ of a graph G are picked independently and uniformly at random, then the partition of $V(G)$ given by the Venn diagram of the neighbourhoods of v_1, \dots, v_ℓ is very likely to have most of its pairs ε -regular. It will not have the property that the classes are all about the same size, but this can be fixed.

4 Sample applications of the Regularity Lemma

Many many modern proofs in graph theory start with the phrase “take an ε -regular partition of G ”. The point is that, once we have taken an ε -regular partition, we know a lot about the structure of the graph already.

One important tool that goes together with the regularity lemma is the counting lemma. We will just state this for triangles here.

Lemma 4.1 (Counting lemma: triangles). *Suppose X, Y, Z are pairwise disjoint sets of vertices in a graph. If all three pairs (X, Y) , (Y, Z) , and (X, Z) are ε -regular with densities d_{XY} , d_{YZ} , and d_{XZ} , then the number of triangles in G with one vertex in each set is*

$$(d_{XY}d_{YZ}d_{XZ} \pm 10\varepsilon)|X||Y||Z|.$$

Proof. Consider the set B of vertices of X with fewer than $(d_{XY} - \varepsilon)|Y|$ neighbours in Y . The set B has size less than $\varepsilon|X|$, since otherwise the pair (B, Y) violates the ε -regularity of (X, Y) . Similarly, less than $\varepsilon|X|$ vertices of X have “too many” neighbours in Y , and the same holds replacing Y with Z . We conclude that at least $(1 - 4\varepsilon)$ vertices of X have $(d_{XY} \pm \varepsilon)|Y|$ neighbours in Y , and $(d_{XZ} \pm \varepsilon)|Z|$ neighbours in Z . By ε -regularity of (Y, Z) , each of these vertices lies in

$$(d_{YZ} \pm \varepsilon) \cdot (d_{XY} \pm \varepsilon)|Y| \cdot (d_{XZ} \pm \varepsilon)|Z|$$

triangles, while the remaining less than $4\varepsilon|X|$ vertices of X lie in at most $|Y||Z|$ triangles each. \square

The full counting lemma provides a similar result for counting copies of any graph. Now one way we can represent the partition of a graph G provided by the regularity lemma is to draw a *reduced graph* $R(G)$ (also called *cluster graph*) whose nodes are the partition classes V_1, \dots, V_k (observe that we are not putting a node for the exceptional set) and whose edges are given a *weight*

³We are ignoring the dependence on k_0 here.

in $[0, 1]$ corresponding to the edge density between the pair of classes. If we define the *triangle density* of G to be the probability that a randomly selected triple of vertices of G form a triangle, then the counting lemma for triangles says that we can approximate the triangle density of G just by looking at $R(G)$.⁴ The full counting lemma then says we can do this for density of any graph H (if $\varepsilon = \varepsilon(H)$ is small enough). So $R(G)$ is a “model” of G whose size is bounded. This turns out to be useful in many proofs.

The counting lemma for triangles implies for example the following surprising result.

Lemma 4.2 (Triangle removal lemma). *For all $\varepsilon > 0$ there is $\delta > 0$ such that any G with at most δn^3 triangles can be made K_3 -free by removing at most εn^2 edges.*

Here is another nice application of the regularity lemma.

Theorem 4.3 (Thomassen 2000; Łuczak 2006). *For each $\eta > 0$ there exists $C = C(\eta)$ with the following property. If G is an n -vertex triangle-free graph whose minimum degree is at least $(\frac{1}{3} + \eta)n$, then G has chromatic number at most C .*

In fact, one can take $C(\eta) = 4$, but this is much more difficult to prove – here we will just show that there is a C . This theorem is best possible in the sense that there is a construction (due to Hajnal, 1973) of triangle-free graphs on n vertices with minimum degree $n/3 - o(n)$ and chromatic number tending to infinity as n tends to infinity.

Proof. Given $\eta > 0$, we set $d = \eta/10$ and $\varepsilon = d^3/10$. We set $k_0 = \varepsilon^{-1}$. The regularity lemma returns a constant $K = K(\varepsilon, k_0)$ and we set $C = C(\eta) = 2^K$.

Now let G be any triangle-free n -vertex graph with minimum degree at least $(\frac{1}{3} + \eta)n$. Let V_0, V_1, \dots, V_k be an ε -regular partition of G with $k_0 \leq k \leq K$ as is guaranteed to exist by the regularity lemma.

Now we define a second partition of $V(G)$ as follows. For each $I \subset [k]$, we let

$$X_I = \left\{ v \in V(G) : \text{we have } |N_G(v) \cap V_i| \geq d|V_i| \text{ if and only if } i \in I, \text{ for } i \in [k] \right\}.$$

This partition has $2^k \leq 2^K = C$ parts, which is independent of n . We claim that all its parts are independent, that is, it witnesses that $\chi(G) \leq C$. We split the proof into two cases.

Case 1, $|I| \geq 2k/3$:

In this case, the set $U = \bigcup_{i \in I} V_i$ has size at least $2n/3 - \varepsilon n$. Thus every vertex of G has at least $(\eta - \varepsilon)n$ neighbours in U , and in particular the average density between pairs from $\{V_i : i \in I\}$ is at least $\eta - 2\varepsilon$ (we “lose” the edges which lie within classes, but there are few of these because the classes are small). Now there are three sorts of pairs from $\{V_i : i \in I\}$ contributing to this average: pairs which are not ε -regular (of which there are few), pairs whose density is smaller than d , and pairs which are ε -regular and of density at least d . The choice of d and ε is such that at least one pair of the latter type occurs (because the average density is as high as $\eta - 2\varepsilon$), say (V_p, V_q) . Let x be any vertex of X_I . Then by definition x has at least $d|V_p|$ neighbours in V_p and at least $d|V_q|$ neighbours in V_q . We conclude by ε -regularity and density of (V_p, V_q) that x is in at least $d^3|V_p||V_q| - \varepsilon|V_p||V_q| > 0$ triangles. This is a contradiction to triangle-freeness of G and we conclude that X_I is actually empty (and so trivially independent).

Case 2, $|I| < 2k/3$:

In this case the set $U = \bigcup_{i \in I} V_i$ has size at most $2n/3$. Since any vertex of X_I has at most $|V_0| \leq \varepsilon n$

⁴In fact we can get the approximation simply from the “weighted triangle density” in $R(G)$. Exercise: Think about how this has to be defined.

neighbours in V_0 , and at most $d|V_i|$ neighbours in any set V_i with $i \notin I$, we conclude that any vertex of X_I has at least $(\frac{1}{3} + \eta - \varepsilon - d)n \geq (\frac{1}{3} + \frac{1}{2}\eta)n$ neighbours in U . But this is more than half of $|U|$, and we conclude that any two vertices in X_I have a common neighbour in U . But then any edge in X_I lies in a triangle. Since G is triangle-free, we conclude that X_I is an independent set in G as desired. \square

The type of proof we saw here is just the start. There are further tools (especially the *blow-up lemma*) that enable more sophisticated applications, including the resolution of a number of what were important open problems in graph theory. Moreover, versions of the Regularity Lemma have been proved for sparse graphs, and for hypergraphs, that enable even more applications. There are also (many) applications outside graph theory. Recall that van der Waerden's Theorem asserts that there exists $n = n(c, r)$ such that in any c -colouring of $[n]$, there is a monochromatic arithmetic progression of length r . Erdős conjectured a stronger statement: if a set S of integers has positive upper density (i.e., $\limsup_{n \rightarrow \infty} |S \cap [n]|/n > 0$), then S contains arbitrarily long arithmetic progressions. This is the theorem proved by Szemerédi in 1975 for which he needed a graph-theoretic lemma, the regularity lemma. In 2012 Szemerédi received the Abel prize, "for his fundamental contributions to discrete mathematics and theoretical computer science, and in recognition of the profound and lasting impact of these contributions on additive number theory and ergodic theory." The regularity lemma played a crucial role in many of these achievements.

5 Exercises

Note: Exercises 2 and 3 are from Bollobás.

1. Fill in the (geometric) details in the proof of Theorem 2.4.
2. Let S be an infinite set of points in the plane. Show that there is an infinite subset A of S such that either no three points of A are on a line, or all points of A are on a line.
3. The Ramsey number $R_k(3)$ is the minimum number n of vertices such that, if the edges of K_n are coloured with k colours, there is always a monochromatic triangle. Show that $R_k(3) \leq k(R_{k-1}(3) - 1) + 2$. [*Hint*: if you do not know the classic proof that $R_2(3) \leq 6$, prove that first.]

Deduce that $R_3(k) \leq \lfloor e \cdot k! \rfloor + 1$.

4. Let B_p be a random bipartite graph, with two vertex classes V_1 and V_2 , each of size n . Each pair of vertices in different classes is joined by an edge with probability p , independently.
 - (a) Show that for all $\varepsilon > 0$, $p > 0$,

$$\mathbb{P}[(V_1, V_2) \text{ is an } \varepsilon\text{-regular pair in } B_{n,p}] \rightarrow 1 \text{ as } n \rightarrow \infty.$$

(You may need some estimates for tails of Binomial random variables.)

- (b) Show that for any bipartite graph H and any fixed $p > 0$,

$$\mathbb{P}[B_{n,p} \text{ contains a copy of } H \text{ as a subgraph}] \rightarrow 1 \text{ as } n \rightarrow \infty.$$

[*Hint*: a crude but straightforward approach starts by taking many disjoint subsets of the vertex set, each of size $|V(H)|$.]

5. Suppose G is a bipartite graph, with vertex classes V_1 and V_2 , each of size n . Suppose also that the maximum degree of G is at most $\varepsilon^2 n$. Show that the pair (V_1, V_2) is ε -regular in G .

6. Let G_n be the following bipartite graph. The vertex set of G_n is $x_1, x_2, \dots, x_n, y_1, y_2, \dots, y_n$. The edges are given by $x_i y_j \in E(G_n)$ if and only if $i < j$.

Fix $\varepsilon > 0$. For each value of n find an explicit ε -regular partition of G_n into at least three and at most (say) $10/\varepsilon$ parts.